# האוניברסיטה העברית בירושלים THE HEBREW UNIVERSITY OF JERUSALEM

### HONESTY OF SIGNALING AND **POLLINATOR ATTRACTION: THE CASE OF FLAG-LIKE BRACTS**

by

### TAMAR KEASAR, GAD POLLAK, RACHEL ARNON, **DAN COHEN and AVI SHMIDA**

Discussion Paper # 438 December 2006

## מרכז לחקר הרציונליות

## **CENTER FOR THE STUDY OF RATIONALITY**

Feldman Building, Givat-Ram, 91904 Jerusalem, Israel PHONE: [972]-2-6584135 FAX: [972]-2-6513681 E-MAIL: ratio@math.huji.ac.il URL: http://www.ratio.huji.ac.il/

## Honesty of signaling and pollinator

## attraction: the case of flag-like bracts

Tamar Keasar<sup>(1\*)</sup>, Gad Pollak<sup>(2)</sup>, Rachel Arnon<sup>(3)</sup>, Dan Cohen<sup>(3)</sup>, Avi Shmida<sup>(3)</sup>

- (1) Dept. of Life Sciences, Achva College, Mobile Post Shikmim 79800, Israel
- (2) Environmental Education program, Kibbutzim College of Education, Tel Aviv, Israel
- (3) Dept. of Evolution, Systematics and Ecology and the Center for Rationality, Hebrew University, Jerusalem 91904, Israel

\* Corresponding author. E-mail: tkeasar@bgu.ac.il; telephone: 972-52-8718860; fax: 972-8-8588057

KEYWORDS: flag-like bract, extra-floral display, pollination ecology, signaling, honeybee, phenology, *Lavandula*, *Salvia* 

RUNNING TITLE: Bee attraction to "flag" displays

#### ABSTRACT

Bracts are nonfloral showy structures associated with inflorescences. They are generally hypothesized to enhance plant reproductive success by attracting pollinating insects. We investigated whether flag-like bracts at the top of inflorescences reliably signal of floral food reward for pollinators in Salvia viridis L. Field and greenhouse data indicate incomplete synchrony between the development of flowers and bracts. Various measures of bract size, however, positively correlate with the number of open flowers on the inflorescence, and with their nectar rewards. Experimental removal of bracts from inflorescences significantly reduced honeybee visitation in the field. We compared these findings with field data on Lavandula stoechas L., another labiate species with flag-like displays. The number of open flowers in L. stoechas cannot be reliably predicted from the presence or size of the bracts. Bract clipping does not significantly reduce honeybee visits in this species. We conjecture that bees learn to orient to those bracts that reliably signal food rewards, and disregard bracts if they provide unreliable signals. Asynchronous development of bracts and floral rewards can reduce the reliability of the signals, and may explain the rarity of flag-like displays in pollination systems. We discuss additional selective forces that may favor bract displays.

#### INTRODUCTION

The evolution of many floral traits is shaped by the foraging behavior of their insect pollinators (Faegri & van-der Pijl, 1979). Pollinators perceive and learn to respond to many features of flowers, such as colors, odors, symmetry, or nectar guides (Giurfa & Lehrer, 2001). These features are considered display cues, which advertise the plant's presence, and / or its quality as a food resource, to potential pollinators. The presence and intensity of the display cues generally correlate positively with nectar and pollen rewards, providing honest advertising (Chittka & Thomson, 2001). Infrequent cases of deception, i.e., non-rewarding flowers with conspicuous displays, presumably involve food mimicry or sexual mimicry (Dafni, 1984; Gigord et al., 2002; Schiestl, 2004).

Colored visual display organs other than flowers are known in many plant species (Heywood 1978; Proctor et al., 1996). They can be divided into two groups: colored organs that encircle the flowers (e.g., *Bouganvillea, Statice* and *Limonium*), and flag-like organs that extend as colorful appendages at the top of inflorescences. Accessory bracts of both groups have been repeatedly hypothesized to provide signals to pollinating insects as to the location of flowers (Muller, 1873; Knuth, 1898; Grant & Grant, 1964; Meeuse & Morris, 1984; Barth, 1985). Bracts were suggested to be very efficient displays, since one large signaling structure, conspicuous from afar, can advertise many flowers on an inflorescence (Faegri & van-der Pijl, 1979; Gottsberger & Hartmann, 1988). Arguably, having more flowers per inflorescence is a simple way to produce large visual displays, which does not require extra-floral display organs. However, such inflorescences may suffer from increased selfing rates through geitonogamy.

Surprisingly, rather few studies attempted to study the advertising role of bracts, and these attempts yielded equivocal results. Bract clipping in a Spanish population of Lavandula stoechas slightly reduced pollinator visitation rates and seed sets, but the effect was generally statistically insignificant (Devesa et al., 1985; Herrera, 1997). The removal of bracts from Mussaenda frondosa L. inflorescences reduced pollination visits by butterflies, but not by bees and birds (Borges et al., 2003). Both studies share a common trend, namely, some reduction in plant reproductive success following bract removal. This trend is consistent with the hypothesis that bracts function as advertising organs. In both studies, however, the effects of bract removal were rather modest, and depended on experimental protocol and types of pollinators observed. The insignificant effects were attributed to small sample sizes and large among-plant variance, which possibly obscured the effects of the bract removal treatment (Herrera, 1997). This interpretation may be limited by lack of information on the correlation between the bract display and the plants' value as food sources. The role of bracts as advertising organs may depend on the reliability of the signal they provide to pollinators as to plant or flower quality, since pollinators may orient mainly to bracts that reliably signal food rewards (Armbruster et al., 2005). Thus, extra-floral displays that dependably predict floral food rewards may be attractive to pollinators, while displays that are poor predictors of floral rewards may be unattractive.

In the present study we explore the correlations between extra-floral displays, pollinator visitation, and floral food reward in two plant species. We focus on the extra-floral displays provided by flag-like bracts. In line with previous studies, we hypothesize that flag-like bracts can function as advertising structures that attract pollinators. We extend this idea and further hypothesize that the effectiveness of this

advertisement depends on its reliability. In other words, extra-floral displays may not provide a reliable foraging signal to pollinators in all plant species or populations. This could occur if display development is not well synchronized with the development and opening of flowers on the inflorescence. In such cases, pollinators should be selected (genetically, or through learning) to disregard the extra-floral signals, and orient to visual or chemical cues originating from the flowers. Pollinators may orient to extra-floral displays, on the other hand, if they honestly advertise the presence of flowers in the inflorescence.

We tested this hypothesis in natural populations of two species with flag-like bract displays in Israel, *Salvia viridis* and *Lavandula stoechas*. We tested whether removal of the flag-like bracts reduced visitation by honeybees. We then assessed the correlation between the extra-floral display and food rewards by studying the plant's flowering phenology in potted plants (for *S. viridis*) or in the field (for *L. stoechas*). We focused on the following questions in each plant species: (1) Are the flag-like bracts effective display cues, i.e., do they attract pollinators? (2) Are the flag-like bracts dependable display cues, i.e., do they reliably signal food rewards to pollinators?

#### METHODS

We sampled bract frequency, monitored flowering phenology, and manipulated extra-floral displays in *Salvia viridis* and *Lavandula stoechas*. Bract frequencies were sampled in natural populations for both species. *S. viridis* phenology was observed in potted plants, while *L. stoechas* phenology was recorded in the field. Manipulations consisted of clipping selected bracts in the field, and recording subsequent pollinator visits. Clipping and observation protocols were not identical for

*S. viridis* and for *L. stoechas*. We therefore describe below the detailed study protocol for each plant species separately.

#### **Study plants**

Salvia viridis (Lamiaceae) is a common annual in Mediterranean and Irano-Turanian grasslands, up to 40 cm tall. It forms dense patches that bloom (in Israel) between March and May. The protandrous flowers are arranged in whorls around the stem, and flowering progresses from the bottom of the inflorescences upwards. The upper lip of the corolla is dark purple, while the lower lip is light purple or white. Corolla length is 15-18 mm (Feinbrun-Dothan 1978; Alon 1990). The species is polymorphic with respect to bracts, since they do not develop in all individuals. In fact, bracted and bractless morphs were previously considered separate species (Davis, 1982; Miekle, 1983). The frequency of the two morphs differs among populations. Flag-like bract clusters, composed of several colorful (purple, pink, or white) leaves, develop at the top of inflorescences. The first bract clusters usually develop on the main inflorescence. In some cases, secondary inflorescences that develop later carry bract clusters as well.

*Lavandula stoechas* L. is a perennial, aromatic, xerophytic shrub of Mediterranean distribution that blooms between January and May (Feinbrun-Dothan, 1978; Herrera, 1993). Flowers are protandrous, and are usually non-selfing and insect-pollinated (Devesa et al., 1985). Flowers have a small (5 mm long), dark-purple tubular corolla inserted into a tubular calyx, and are aggregated in heads. These are composed of tightly packed groups of flowers, attached to a central common axis. In preliminary observations, we determined that  $15.0\pm 8.3$  (mean  $\pm$  SD, n=58) flowers bloom simultaneously per inflorescence, and that mean flower longevity is  $4.61\pm 1.54$ 

(SD) d (n=70). The mean number of inflorescences per shrub is  $117.5\pm87.9$  (mean  $\pm$  SD, n=30) at peak blooming. Many inflorescences are terminated by a tuft of 4-6 pink bracts. Mean inflorescence length is  $22.0\pm5.9$  (SD, n=100) mm, and mean length of the bract cluster is  $7.0\pm5.9$  (SD, n=85) mm. Thus, bracts account for about a fourth of total inflorescence size. These measurements agree well with data obtained for *L*. *stoechas* populations in Spain (Devesa 1985; Gottsberger & Hartmann 1988).

#### **Study sites**

The main field site for *S. viridis* was the Ruhama nature reserve, in the southwest of Israel. The area is characterized by phrygana vegetation dominated by *Thymelea hirsuta*, *Hyparrhenia hirta*, and *Coridothymus capitatus* growing on loess soil. Honeybees and solitary bees dominate the pollinator fauna. *S. viridis* forms large patches, made conspicuous by purple bracts, in the reserve.

*L. stoechas* was studied at an open natural vegetation area near Harutzim in central Israel. The soil at the study site is red sandy loam, and the vegetation is a mosaic of annual pasture patches and tall phrygana. The perennial vegetation is dominated by *Cistus salviifolius* and *Cistus creticus*, accompanied by *Calycotome villosa* and *Thymelaea hirsuta*. Honeybee hives were located 200 m off the research site. Honeybees accounted for more than 99% of the pollinator visits observed. The remaining pollinators were solitary bees (*Eucera* sp., *Nomada* sp.), wasps (*Polystes* sp.), beetles (*Oxythyrea*), moths (*Zygaena*), and two unidentified heteropterans.

#### Field sampling of bract frequency

We sampled the prevalence of bracts on flowering *S. viridis* plants in four natural populations of *S. viridis* in northern (Brosh), central (Pura) and southern

(Dorot, Ruhama) Israel in 2000 and 2001. For the Ruhama population, we also recorded the frequency of inflorescences that carried bracts without flowers. The prevalence of bracts in *L. stoechas* was sampled near Harutzim in central Israel in 1990.

#### Phenological observations – S. viridis

We used field-collected *S. viridis* seeds to produce 51 potted plants, grown under greenhouse conditions, in 2000. Seeds produced by these plants were used to grow 16 additional shrubs in 2001. We recorded the lengths of all bracts, the number of leaves per bract cluster, and the number of bracts per plant as measures of display intensity. We recorded the total number of flowers per plant, and determined nectar volumes in samples of flowers as indicators of the plant's food reward. We measured nectar volumes in samples of six flowers per inflorescence, taken from the bottom (two flowers), mid-height (two flowers), and top (two flowers) of the inflorescence. All parameters were recorded at three-day intervals during the plants' two-month flowering period. In 2001, we recorded the presence of bracts and of flowers in 16 plants once a day, to obtain precise data on the degree of synchrony between them.

#### Phenological observations – L. stoechas

We conducted weekly counts of the number of blooming inflorescences in a sample of thirty shrubs at the Harutzim study site to characterize the time course of blooming. We counted the number of open flowers in 60 inflorescences three times a week throughout the blooming period.

We registered the dates of appearance and wilting of the bracts in 100 marked inflorescences. One inflorescence broke during the study period. Data from the

remaining 99 inflorescences were used for analysis. We recorded blooming dates for these inflorescences, i.e., the dates of opening of the first flower, and of wilting of the last flower of the inflorescence. These records provide information on the extent of synchrony between flower and bract cluster development.

#### Bract manipulation experiments – S. viridis

Experiments were conducted on three days in the spring of 2001. In all experiments, we clipped all bracts from S. viridis inflorescences. We recorded the number of arrivals of unmarked honeybees to manipulated inflorescences and to an equal number of intact control inflorescences. Since bees generally visited more than one flower per inflorescence, we also recorded the total number of visits to flowers on clipped and unclipped inflorescences. We followed each bee until it left our experimental patches. We used the total number of visit sequences as an estimate to the number of visiting individuals. This is probably an overestimate, since some individuals likely visited the experimental patches more than once during the observation periods. Visits were recorded by two observers immediately after bract cluster removal, simultaneously for manipulated and control inflorescences. We used three bract cluster removal treatments, as detailed in Table 1: (a) "Patch" treatment, where we clipped bract clusters from a whole patch; (b) "Binary" treatment that involved removal of one half of the bract clusters within a patch at random locations; and (c) "Split" treatment, in which we clipped all bract clusters from one half of a patch, and left its other half intact. In one of the "patch" treatments we clipped and reattached all bract clusters in the control patch, while bract clusters in the manipulated patch were clipped and removed (clipped control, Table 1). This was

done to control for the possibility that odors emanating from clipped inflorescences deterred pollinators.

We determined nectar volumes in a sample of flowers from manipulated and control inflorescences in eight replicates, to test for possible effects of clipping on floral reward levels. We used 1- $\mu$ l microcapillaries for nectar sampling. Sample sizes for nectar measurements are provided in Table 1.

#### TABLE 1

#### **Bract manipulation experiments** – *L. stoechas*

Experiments were conducted on nine days in the spring of 1990. As in *S*. *viridis*, we compared pollinator activity on intact infloresences vs. inflorescences from which bract clusters had been experimentally removed. We created two bract cluster removal treatments. In the "Plant" treatment, we clipped bract clusters at the bud stage from all inflorescences of treated plants, creating plants that bore no bracts throughout the blooming season. In the "Inflorescence" treatment, we clipped bract clusters from haphazardly determined inflorescences just before observation sessions. This created plants that contained mosaics of bracted and bractless inflorescences (Table 2). We set up these treatments to control for the possibilities that (a) pollinators can memorize the locations of bracts (and orient to these locations) even after the bracts are no longer there, and (b) a large concentration of bract clusters increases pollinator attraction. If these possibilities are valid, then pollinators are expected to discriminate against clipped inflorescences in the "Plant" treatment, but not in the "Inflorescence"

treatment. We recorded the number of pollinator arrivals to manipulated and control inflorescences simultaneously in ten-minute observation sessions.

#### TABLE 2

#### Data analysis

We calculated the proportion of sampled inflorescences that bore both flowers and bracts, flowers without bracts, and bracts without flowers, in both study species. In *S. viridis*, we used linear regressions to relate bract display parameters (bract cluster length, no. leaves per bract cluster, no. bract clusters per plant) to reward parameters (number of flowers, mean nectar volume per flower). We used each plant's measurements of display and reward on the day of maximal bract cluster length for analysis. Thus, each plant in the sample contributed a single data point to the regression. This analysis was not performed for *L. stoechas*, since bract clusters did not change in size during the blooming period.

In the bract removal experiments, we scored the proportion of replicates that had more bee visits to intact inflorescences than to clipped inflorescences. We tested the hypothesis that this proportion was higher than 0.5 using one-way sign tests. We employed Wilcoxon paired-sample tests to examine whether mean parameters of bee visits were higher in control inflorescences than in their manipulated counterparts.

#### RESULTS

#### S. viridis

#### The prevalence of bracts, and bract-flower synchrony

Blooming inflorescences bore bracts in almost all plants sampled in field populations, and in potted plants in the greenhouse (Table 3). In the Ruhama population, 23% of the sampled inflorescences bore bracts but no open flowers, suggesting incomplete synchrony between flowering and the development of the potential advertising signal. In potted greenhouse plants, bracts developed  $9.47 \pm 1.70$ d (mean $\pm$ SE, n=17) before the blooming of the first flowers, and wilted  $2.00 \pm 1.29$  d after the last flowers of the inflorescence. Bracts preceded flowers in 94% of this 17plant sample, and remained on the plant after the flowers had wilted in 29% of the plants in the sample.

#### TABLE 3

#### **Bract cluster size and reward parameters**

The number of flowers first increased, then decreased during the blooming period. Plants developed secondary inflorescences during the flowering period. Since secondary inflorescences often bore bracts, the number of bract clusters per plant changed along the flowering period as well. Bract clusters increased in length through the addition of leaves, and decreased in length when some of these leaves wilted. A typical time course for the changes in flower number, bract cluster number, bract cluster length, and number of leaves per bract cluster is provided in Fig. 1.

Bract cluster length was significantly and positively correlated with the number of flowers per inflorescence (n=65,  $r^2$ =0.31, p<0.0001) and nectar volume per

flower (n=65,  $r^2$ =0.31, p<0.0001). Flower number and nectar content also correlated positively with the number of bract clusters (n=65,  $r^2$ =0.14, p<0.0001 and  $r^2$ =0.19, p=0.002, respectively) and with the number of leaves per bract cluster (n=65,  $r^2$ =0.13, p=0.002 and 0.19, p=0.001, respectively).

FIG 1

#### **Bract manipulation experiments**

Honeybees visited fewer flowers in manipulated inflorescences than in untreated control plants in ten out of twelve experiments (all three treatments were pooled). In these ten experiments, the number of arrivals at control inflorescences was also higher than at bractless inflorescences (Table 4). The occurrence of the same pattern in ten out of twelve cases is unlikely to result from a random process (sign test, n=12, Z=2.02, P=0.04). The number of bees in control patches was higher than in manipulated patches in eight out of eleven replicates (the number of bees was not recorded in replicate no. 3), i.e., not more frequently than expected at random. We pooled the twelve replicates from different treatments, and calculated the difference between treatment and control in the number of arrivals at inflorescences, visits to flowers, and number of bees. The significance of the difference was 0.058 (Z=1.57, n=12) for arrival at inflorescences, 0.050 (Z=1.65, n=12) for visits to flowers, and 0.046 (Z=1.69, n=11) for number of bees (one-tailed Wilcoxon matched-pairs tests). Nectar volumes in clipped inflorescences did not differ significantly from nectar volumes in control inflorescences (Mann-Whitney U-test, p=0.32), suggesting that the clipping treatment did not inhibit nectar production. Bees preferred the "clipped

control" inflorescences over manipulated inflorescences in replicate no.1. This suggests that the injury inflicted on the plants by clipping did not, by itself, repel pollinators.

#### TABLE 4

#### L. stoechas

#### The prevalence of bracts, and bract-flower synchrony

70.4% of 281 blooming inflorescences sampled on three dates bore bracts. Eleven out of 99 inflorescences followed throughout the blooming season did not develop any bract. In the remaining inflorescences, bracts were maintained for  $33.58\pm13.91$ (mean $\pm$ SD) d, while flowering lasted  $46.56\pm6.34$  d. Bracts appeared  $4.83\pm3.72$  d before the onset of flowering, and wilted  $18.05\pm13.91$  before flowering ended. Flowers and bracts overlapped for  $27.59\pm11.73$  d, that is, during  $59.59\pm25.88\%$  of the duration of blooming. Taking into account the fact that bracts were totally absent in 11% of the inflorescences (i.e., zero overlap between bracts and flowers), the probability that a blooming inflorescence would carry a bract was 0.54. The probability that a bract would signal a blooming inflorescence was 0.86.

#### Bract size and the number of open flowers

The blooming period of our study population extended from early January to early May, and peaked between mid-March and mid-April. The blooming span of individual shrubs was 90.83±17.19 (Mean±SD, n=30). We constructed a blooming diagram for each of the sixty inflorescences surveyed during this period. Blooming occurred in each inflorescence in 3-4 waves. The number of open flowers typically peaked four days after the onset of blooming. The second and third peaks occurred

after 12-18 and 25-35 days, respectively. The fourth peak, which was much less pronounced (and at times missing altogether), occurred at least 40 days after the onset of flowering. Bracts were usually present in the inflorescence during the first 1-2 blooming peaks, but not during the third and fourth peaks. Bract cluster size remained unchanged during the whole display period.

#### **Bract manipulation experiments**

Pollinators landed on intact inflorescences more frequently than on bractless inflorescences in 9 out of 14 observations. This frequency does not significantly differ from 0.5 (sign test, n=14, P=0.212). The number of visits to intact inflorescences was higher than to clipped inflorescences in 11 out of 17 observations. This preference is not statistically significant either (sign test, n=17, P=0.166). The mean numbers of pollinator arrivals and flower visits were not significantly affected by bract cluster removal (Wilcoxon matched-pairs tests, n=14, Z=1.04, p=0.30 for number of arrivals, n=17, Z=0.87, p=0.38 for number of visits).

#### DISCUSSION

The main manipulation in the present study involved the removal of flag-like bracts from the inflorescences of two plant species of the mint family. Following the manipulation, honeybees made fewer visits to *S. viridis* plants that lacked flag-like bracts than to control plants. This result supports the hypothesis that terminal clusters of colored bracts function as advertising organs that attract pollinators. In this experiment, we did not control for possible memorization of bract locations by pollinators. Thus, some of the bees' visits to clipped inflorescences may reflect their memory of patches that had borne bracts in the past. A similar reduction in pollinator visits was obtained by petal removal from flowers in plant species that lack bracts

(Bell, 1985, Kudoh & Whigham, 1998). We obtained different results for *L. stoechas*: honeybees did not significantly prefer control inflorescences over clipped ones. This finding concurs with similar previous manipulations on *L. stoechas* (Devesa et al., 1985, Herrera, 1997), but run counter to the regnant paradigm that pollinators prefer flagged inflorescences (Faegri & van-der Pijl, 1979; Barth, 1985; Gottsberger & Hartmann, 1988; Proctor et al., 1996).

The two study species also differed in the correlation between bract display and food reward, i.e., in the reliability of bracts as advertising cues. Bracts provide more reliable signals of food rewards in *S. viridis* than in *L. stoechas* in two respects: (1) In *S. viridis*, bracts may provide false positive signals of reward (bracts exist in the absence of open flowers), but false negative signals (open flowers with no bracts) are very rare. In *L. stoechas*, bracts are frequently associated with both false positive and false negative signals. (2) In *S. viridis*, various measures of display size (number of flag-like bract clusters, number of leaves per bract cluster, bract cluster length) are consistently and positively correlated with reward parameters. This is not the case in *L. stoechas*, where the number of open flowers, but not bract cluster size, fluctuates in each inflorescence during the blooming season.

Our data thus show that *Salvia* bracts are both more effective and more reliable display cues than *Lavandula* bracts as signals that indicate nectar rewards in the inflorescences below them. It is tempting to posit a cause-and-effect relationship between the reliability and the effectiveness of the flag-like displays: pollinators may learn to orient to *Salvia* bracts, but may also learn to disregard *Lavandula* bracts as dishonest advertising signals. Alternatively, *S. viridis* bracts may be more attractive to pollinators because they provide a larger visual stimulus, relative to the inflorescence. *S. viridis* bract signals may exert a greater effect on pollinators' patch choices because

they are detectable from a greater distance (Vaknin et al., 1996). Pollinators learn display cues mainly on their way to the food source. This learning requires at least three seconds of exposure to the display stimulus (Menzel, 1985). As detection distance increases, the pollinator spends more time *en route*, allowing more time for learning the display cue. This mechanism suggests that large bracts (as in *S. viridis*) may affect pollinator choices more strongly than small bracts (as in *L. stoechas*).

Our results do not point to any evolutionary advantage to the asynchronous development of flag-like bracts and flowers in L. stoechas. Unlike other cases of deception in plant-pollinator systems (e.g., deceptive orchids, Dafni, 1984), no pollination benefit is expected for plants that attract pollinators by carrying flag-like bracts, but do not bloom. Moreover, the maintenance of bracts before blooming may reduce the total amount of resources available for reproduction, and may therefore be maladaptive. A possible interpretation is that the asynchrony between bracts and flowers reflects a developmental, non-adaptive constraint (Herrera, 2001): from a pollination point of view, flag-like inflorescences can be viewed as equivalent to flowers, because they contain display, reward and sex structures. Unlike flowers, however, the display structure is situated at a distance from the sex and reward organs and is related to the whole inflorescence, rather than to a single flower. This feature may pose a constraint on synchronized development of flowers and displays. Indeed, bract displays remain conspicuous in several species (e.g. Salvia sclarea, Bougavillea spp.) long after the flowers associated with them have finished blooming. Constraints on the synchronized development of bracts and flowers are likely to limit the selective advantage of flag-like bracts. This may explain why flag-like bracts are such a rare phenomenon in plants: out of ca. 4500 genera in the Mediterranean flora only four genera possess species with flag-like bracts: Muscari, Leopoldia, Lavandula and

*Salvia*. While the flag-like displays of *Salvia* and *Lavandula* are composed of bract clusters, the displays of *Muscari* and *Leopoldia* are formed by aborted, transformed flowers. Regardless of their morphological origin, the displays are colored and much more conspicuous than the flowers within the same inflorescence.

An alternative explanation for the asynchrony of flowers and flag-like bracts is that bracts have evolved for a function other than pollinator attraction, such as defense from radiation damage, drought, or herbivory (Galen & Cuba, 2001; Armbruster, 2002). According to this hypothesis, the presence of anthocyanins in vegetative organs, such as bracts, is selected for reasons unrelated to pollination. If the main function of bracts is indeed chemical defense from herbivores, then they should be selected to appear during the period of maximal grazing pressure. Such selection could decouple bract development from flowering, if the time of maximal herbivory does not coincide with blooming (but see Herrera, 1993, for temporal patterns of herbivory on *L. stoechas*).

The similarity in pollinator visitation rates to bract-bearing and bractless inflorescences of *L. stoechas* begs the question of whether any pollination-related selective pressures favor bract development in this species. The following selective advantages have been proposed: (a) Flag-like bracts may increase *L. stoechas* detectability to pollinators mainly in low-density, establishing populations (Herrera, 1997); (b) Flag-like bracts convey information on the location of *L. stoechas* shrubs mainly to young naïve foragers on their very first flights. As these pollinators gain experience, they learn other cues associated with the plants, such as their location (Wehner & Menzel, 1990), thereby reducing their reliance on the bract display; (c) Flag-like bracts may attract pollinators from long distances to the general area of the flowering patch, but insects' choice of specific inflorescences at short distances is

guided by different cues. In other words, bracts may function as "detective cues" that advertise a plant's location rather than as "selective cues" that advertise its quality (Lewis & Lipani, 1990; Cohen & Shmida, 1993); (d) Flag-like bracts may provide plants with a mechanism to discriminate between potential pollinators, and to evade the less efficient ones (Proctor et al., 1996). Preliminary observations in four species of plants with flag-like bracts indicate that flies direct a higher proportion of their flights to bracts (rather than flowers) as compared to bees (Table 5). Similarly, removal of bracts from *Mussaenda frondosa* inflorescences reduced pollination visits by butterflies, but not by bees and birds (Borges et al., 2003). Bees are considered more efficient pollinators than flies and butterflies, since they fly longer distances and are more flower-constant (Waser, 1983). The presence of flag-like bracts may thus increase the probability of pollination by bees (Menzel & Shmida, 1993). We were not able to compare bee vs. fly attraction to bracts in the present study, as almost all observed pollinators were honeybees. We suggest that this issue deserves further study in field sites that contain several pollinator groups.

#### ACKNOWLEDGMENTS

We thank Yuval Sapir, Revital Kashi, Noam Raanan and Noam Bar-Shai for their help with field observations. Haim Efrat kindly provided honeybee colonies for preliminary observations. Scott Armbruster commented on an earlier version of the manuscript.

#### REFERENCES

Alon, A. 1990. Israel's Fauna and Flora: Flowering Plants. Peli, Ramat Gan (in Hebrew).

Armbruster, W.S. 2002. Can indirect selection and genetic context contribute to trait diversification? A transition-probability study of blossom-colour evolution in two genera. J. Evol. Biol. 15: 468-486.

Armbruster, W. S ,.Antonsen, L., Pélabon, C. 2005. Phenotypic selection on blossoms in an African *Dalechampia* :honest signaling, visitation rates, and cross-pollination success. Ecology 86: 3323–3333.

Barth, F.G.1985. Insects and flowers: The biology of a partnership. Princeton University Press. Princeton, N.J. 297 pp.

Bell, G. 1985. On the function of flowers. Proc. Roy. Soc. Lond. B 224: 223-265.

Borges, R.M., Gowda, V., Zacharias, M. 2003. Butterfly pollination and high-contrast visual signals in a low-density distylous plant. Oecologia 136: 571-573.

Chittka, L. and Thomson, J.D. (eds.) 2001 Cognitive Ecology of Pollination - Animal Behavior and Floral Evolution. Cambridge University Press., Cambridge.

Cohen, D., Shmida, A. 1993. The evolution of flower display and reward. Evol. Biol. 27: 197-243.

Dafni, A. 1984. Mimicry and deception in pollination. Ann. Rev. Ecol. Syst. 15: 259-278.

Davis, P. E. 1982 . Flora of Turkey and the East Agean Islands. Vol 7. Univ. Press, Edinburgh, Edinburgh .

Devesa, T. A., Arroyo, J., Herrera, J. 1985. Contribution to the knowledge of the floral biology of the genus *Lavandula*. Anales Jard. Bot. Madrid 42: 165-186 (in Spanish).

Faegri, K., van der Pijl, L. 1979. The principles of pollination ecology. Pergamon Press, London.

Feinbrun-Dothan N. 1978 *Flora Palaestina*. *Part 3*. Israel Academy of Sciences and Humanities, Jerusalem.

Galen, C., Cuba, J. 2001. Down the tube: pollinators, predators, and the evolution of flower shape in the alpine skypilot, Polemonium viscosum. Evolution 55: 1963-1971. Giurfa, M., Lehrer, M. 2001. Honey bee vision and floral displays: from detection to close-up recognition. In: Chittka L, Thomson JD (Eds.). Cognitive ecology of pollination: animal behavior and floral evolution. Cambridge University Press, Cambridge. pp. 61-82.

Gigord, L.D.B., Macnair, M.R., Stritesky, M., Smithson, A. 2002. The potential for floral mimicry in rewardless orchids: An experimental study. Proc. Roy. Soc. Lond. B 269: 1389-1395.

Gottsberger, G., Hartmann, U. 1988. Flag-blossom and long-lasting flowers: A strategy for effective pollinator attraction in the Mediterranean region and in Atlantic Potugal. Lagascalia, 15: 635-641.

Grant, K.A., Grant, V. 1964. Mechanical isolation of *Salvia apiana* and *Salvia mellifera* (Labiatae). Evolution 18: 196-212.

Herrera, J. 1993. Blooming times of individual inflorescences and plants as determinants of flower and fruit predation in *Lavandula stoechas* (Lamiaceae). Acta Oecologica 14: 867-874.

Herrera, J. 1997. The role of colored accessory bracts in the reproductive biology of *Lavandula stoechas*. Ecology 78: 494-504.

Herrera, C.M. 2001. Deconstructing a floral phenotype: Do pollinators select for corolla integration in *Lavandula latifolia*? J. Evol. Biol. 14: 574-584.

Heywood, V.H., 1978. Flowering plants of the world. Oxford University Press, Oxford.

Knuth, P. 1988. Handbuch der Blutenbilogie. Verlag von Wilhelm Ingelmann, Leipzig. Vol 5.

Kudoh, H., Whigham, D.F. 1998. The effect of petal size manipulation on pollinator /seed-predator mediated female reproductive success of *Hibiscus moscheutos*. Oecologia 117: 70-79.

Lewis, A. C., Lipani, G. A. 1990. Learning and flower use in butterflies:

Hypothesis from honey-bees. In Bernays, E. A. (ed.) Insect-Plant Interactions.

CRC Press, Boca Raton, Fl., pp. 95-110.

Meeuse, B., Morris, S. 1984. The sex life of flowers. Faber & Faber, London.

Miekle, R.D. 1985. Flora of Cyprus, vol. 2. Royal Botanic Gardens, Kew.

Muller, H. 1883. The fertilization of flowers. MacMillan Press & Co., London.

Proctor, M., Yeo, P., Lack, A., 1996. The Natural History of Pollination. Harper Collins, Hampshire, UK.

Schiestl, F.P. 2004. Floral evolution and pollinator mate choice in a sexually deceptive orchid. J. Evol. Biol. 17: 67-75.

Vaknin, Y., Yom-Tov, Y., Eisikowitch, D., 1996. Flowering seasonality and flower characteristics of *Loranthus acaciae* Zucc. (*Loranthaceae*): implications for advertisement and bird-pollination. Sex. Plant Rep. 9: 279 - 285.

| Pattern of bract | Replicate | No.            | Duration of | No. flowers for   | No. flowers for   |
|------------------|-----------|----------------|-------------|-------------------|-------------------|
| removal          |           | inflorescences | observation | nectar sampling - | nectar sampling - |
|                  |           | observed       | (min.)      | manipulated       | Control           |
|                  |           | (manipulated+  |             |                   |                   |
|                  |           | control)       |             |                   |                   |
| Patch, clipped   | 1         | 180            | 180         | 32                | 37                |
| control          |           |                |             |                   |                   |
| Patch            | 2         | 170            | 110         | 29                | 33                |
| Patch            | 3         | 120            | 120         | 19                | 24                |
| Patch            | 4         | 15             | 50          | -                 | -                 |
| Patch            | 5         | 15             | 140         | -                 | -                 |
| Patch            | 6         | 55             | 45          | -                 | -                 |
| Binary           | 1         | 50             | 30          | 25                | 22                |
| Binary           | 2         | 122            | 60          | 11                | 17                |
| Binary           | 3         | 120            | 50          | 35                | 24                |
| Split            | 1         | 50             | 30          | 19                | 25                |
| Split            | 2         | 100            | 60          | 34                | 31                |
| Split            | 3         | 120            | 50          | -                 | -                 |

Table 1: Details of bract manipulations and nectar sampling in the Salvia experiments.

Table 2: Pollinator activity in *L. stoechas* bract manipulation experiments. "plant" and "Inflorescence" treatments differ in the protocol of bract clipping (see methods). We report the mean number of pollinator arrivals, and the mean number of flowers visited per inflorescence in a standard 10- min observation period. NR – not recorded. Cases of lower pollinator activity in manipulated inflorescences than in controls are marked in bold. All pollinators were honeybees, except in replicates 2, 3 in treatment "inflorescence" (some visits by *Eucera* sp.) and replicate 8 in treatment "inflorescence" (one visit by *Oxythyrea* sp.)

| Replicate | No. inflorescences |         | No. pollinator arrivals |         | No. flowers visited |         |
|-----------|--------------------|---------|-------------------------|---------|---------------------|---------|
|           | observed           |         |                         |         |                     |         |
|           | Clipped            | Control | Clipped                 | Control | Clipped             | Control |
| 1         | 86                 | 161     | NR                      | NR      | .558                | .509    |
| 2         | 56                 | 45      | NR                      | NR      | 1.050               | .490    |
| 3         | 55                 | 68      | NR                      | NR      | .600                | 1.260   |
| 4         | 67                 | 42      | NR                      | NR      | .630                | 1.190   |
| 5         | 95                 | 28      | .042                    | .178    | .378                | .714    |
| 6         | 87                 | 57      | .149                    | .333    | .632                | 1.400   |
| 7         | 53                 | 29      | .472                    | .724    | 1.940               | 3.370   |

a. Treatment "Plant"

### b. Treatment "Inflorescence"

| Replicate | No. inflorescences |         | No. pollina | No. pollinator arrivals |         | No. flowers visited |  |
|-----------|--------------------|---------|-------------|-------------------------|---------|---------------------|--|
|           | observed           |         |             |                         |         |                     |  |
|           | Clipped            | Control | Clipped     | Control                 | Clipped | Control             |  |
| 1         | 71                 | 56      | .098        | .107                    | NR      | .839                |  |
| 2         | 26                 | 28      | .385        | .357                    | 1.269   | 1.428               |  |
| 3         | 26                 | 28      | .192        | .285                    | .423    | 1.178               |  |
| 4         | 26                 | 28      | .692        | .321                    | 2.730   | 1.320               |  |
| 5         | 29                 | 26      | .172        | .423                    | 1.000   | 1.460               |  |
| 6         | 43                 | 28      | .023        | .214                    | .093    | .928                |  |
| 7         | 32                 | 40      | .063        | .125                    | .312    | 1.050               |  |
| 8         | 73                 | 124     | .041        | .048                    | .465    | .177                |  |
| 9         | 73                 | 107     | .109        | .037                    | 1.315   | .336                |  |
| 10        | 49                 | 29      | .082        | .068                    | .489    | .621                |  |
| 11        | 49                 | 29      | .184        | .000                    | .979    | .000                |  |

| Population | No. plants sampled | Proportion of blooming     |
|------------|--------------------|----------------------------|
|            |                    | inflorescences with bracts |
| Brosh      | 150                | 0.99                       |
| Pura       | 150                | 1                          |
| Dorot      | 150                | 1                          |
| Ruhama     | 245                | 0.89                       |
| Greenhouse | 67                 | 0.98                       |

## Table 3: Proportion of blooming S. viridis inflorescences that bore bracts

Table 4: Pollinator activity in S. viridis bract manipulation experiments. Cases of lower pollinator activity in manipulated inflorescences

than in controls are marked in bold.

| Pattern of bract removal | Replicate | No. flowers visited |             | No. inflorescences visited |             | No. bees     |              |
|--------------------------|-----------|---------------------|-------------|----------------------------|-------------|--------------|--------------|
|                          |           | Control             | Manipulated | Control                    | Manipulated | Control      | Manipulated  |
| Patch, clipped control   | 1         | 499                 | 326         | 384                        | 157         | 19           | 18           |
| Patch                    | 2         | 335                 | 185         | 239                        | 143         | 10           | 11           |
| Patch                    | 3         | 295                 | 467         | 133                        | 305         | Not recorded | Not recorded |
| Patch                    | 4         | 0                   | 138         | 0                          | 82          | 0            | 7            |
| Patch                    | 5         | 253                 | 113         | 161                        | 100         | 44           | 45           |
| Patch                    | 6         | 112                 | 60          | 118                        | 50          | 16           | 12           |
| Binary                   | 1         | 102                 | 47          | 55                         | 27          | 5            | 1            |
| Binary                   | 2         | 300                 | 272         | 211                        | 201         | 13           | 9            |
| Binary                   | 3         | 135                 | 117         | 109                        | 80          | 5            | 2            |
| Split                    | 1         | 442                 | 276         | 261                        | 166         | 20           | 6            |
| Split                    | 2         | 112                 | 22          | 71                         | 18          | 5            | 2            |
| Split                    | 3         | 25                  | 16          | 18                         | 11          | 3            | 0            |

Table 5: Observations on bee and fly behavior on inflorescences with flag-like bracts. One hundred pollinator visitation sequences were observed for each plant species. For each visitation sequence, we noted the type of pollinator (fly or anthophorid bee), and whether it included the flag-like bracts, the flowers, or both.

| Plant species  | Inflorescences | Pollinator type  | No. of visitation sequences to |         |            |  |
|----------------|----------------|------------------|--------------------------------|---------|------------|--|
|                | observed       |                  |                                |         |            |  |
|                |                |                  | Flag-like                      | Flowers | Flowers    |  |
|                |                |                  | Bracts                         | only    | and bracts |  |
|                |                |                  | only                           |         |            |  |
| Leopoldia      | 2              | Flies            | 12                             | 1       | 3          |  |
| comosa         |                |                  |                                |         |            |  |
|                |                | Anthophorid bees | 0                              | 81      | 3          |  |
| Salvia viridis | 1              | Flies            | 5                              | 1       | 0          |  |
|                |                | Anthophorid bees | 0                              | 94      | 0          |  |
| Muscari        | 3              | Flies            | 33                             | 1       | 15         |  |
| commutatum     |                |                  |                                |         |            |  |
|                |                | Anthophorid bees | 0                              | 48      | 2          |  |
| Eremurus       | 1              | Flies            | 4                              | 0       | 1          |  |
| spectabilis    |                |                  |                                |         |            |  |
|                |                | Anthophorid bees | 0                              | 95      | 0          |  |

#### FIGURE CAPTIONS

Fig. 1: Representative time course of blooming and bract cluster development in a *S*. *viridis* inflorescence. Filled squares – number of flowers, empty squares – number of bracts, filled triangles – number of bract clusters, empty triangles – bract cluster length (cm). Best-fit curves are based on 6<sup>th</sup>-order polynomials.

Fig. 1

